

THE BRAND MOMS TURN TO FIRST*




*When they first treat their children's upper respiratory allergies.

AGE	DOSE
6-11 years	Take 5 to 10 mL every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.

AGE	DOSE
12+ years	Take 10 mL every 4 hours. Do not take more than 6 doses in 24 hours.
6-11 years	Take 5 mL every 4 hours. Do not take more than 6 doses in 24 hours.
4-5 years	Do not use unless directed by a doctor.
Under 4 years	Do not use.

AGE	DOSE
12+ years	2 to 4 chewable tablets (25 mg to 50 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
6-11 years	1 to 2 chewable tablets (12.5 mg to 25 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.



AVAILABLE IN
DYE, SUGAR,
& ALCOHOL
FREE

**Children's
BENADRYL®
Allergy Liquid[‡]**

Use products only as directed.



**Children's
BENADRYL®
Allergy Plus
Congestion^{‡†}**

Use product only as directed.



**Children's
BENADRYL®
Chewables[‡]**

Use product only as directed.



Help your children stop itching with the cooling, soothing itch relief of **BENADRYL® Anti-Itch Gel For Kids.**

For additional information on allergy management, talk to your pediatrician.

†Relieves runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.

‡Use only with dosing cup provided with product.

mL = milliliter