

Children 6 months to 11 years

MOTRIN® Pediatric Products

DOSE: Every 6-8 hours as needed

Do NOT give more than 4 doses in 24 hours

If possible, use weight to dose; otherwise, use age.



Infants' MOTRIN® Concentrated Drops

Active ingredient: ibuprofen (NSAID)* 50 mg (in each 1.25 mL)

Available in: Berry Dye-Free Berry

WEIGHT	12-17 lbs	18-23 lbs
AGE	6-11 months	12-23 months
DOSE	1.25 mL 	1.875 mL

Use product only as directed.



Children's MOTRIN® Oral Suspension

Active ingredient: ibuprofen (NSAID)* 100 mg (in each 5 mL)

Available in: Grape Berry Dye-Free Berry Bubble Gum

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	5 mL 	7.5 mL 	10 mL 	12.5 mL 	15 mL

Use product only as directed.

mL = milliliter



Children's MOTRIN® Chewables

Active ingredient: ibuprofen (NSAID)* 100 mg (in each chewable tablet)

Available in: Grape Dye-Free Grape

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	1 tablet 	1½ tablets 	2 tablets 	2½ tablets 	3 tablets

Use product only as directed.

Important instructions for proper use: Always read and follow the label on all MOTRIN® products. Use only the dosing device (dosing syringe or dosing cup) that came with the product. Do **NOT** use any other dosing device.

*Nonsteroidal anti-inflammatory drug.

Adults and children 12 years and older

MOTRIN® Adult Products*



MOTRIN® IB Liquid Gels

Active ingredient: ibuprofen (NSAID)† 200 mg (in each capsule)

Use product only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY	1 capsule every 4-6 hours while symptoms last; if pain or fever does not respond to 1 capsule, 2 capsules may be used
MAXIMUM LABELED DOSE	Not to exceed 6 capsules in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 1200 mg</i>



MOTRIN® IB Caplets

Active ingredient: ibuprofen (NSAID)† 200 mg (in each caplet)

Use product only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY	1 caplet every 4-6 hours while symptoms last; if pain or fever does not respond to 1 caplet, 2 caplets may be used
MAXIMUM LABELED DOSE	Not to exceed 6 caplets in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 1200 mg</i>

Important instructions for proper use: Always read and follow the label on all MOTRIN® products.

†Nonsteroidal anti-inflammatory drug.

*This is not a complete list of MOTRIN® adult products.

Find additional practice and patient resources at TylenolProfessional.com and JnJPediatrics.com